```
Factorizing Cubic Equations Of The Form Ax^3 + Bx^2 + Cx + D = 0
```

document.write(a, ", ", b, ", ", c, ", ",

}

}

}

</script> </body> </html>

" = ", a*b + a*c + b*c, "
");

Written by Adi Cox 4th May 2013

```
Where a, b, c are the integer solutions to the cubic equation if
and only if A = 1.
B = a+b+c
C = ab+ac+bc
D = abc
The html file below finds the value of C in the cubic equation
for the values B and D that are entered.
<!DOCTYPE HTML PUBLIC " - //W3C//DTD XHTML 1.0 Transitional//EN"</pre>
"http://www.w3.org/TR/xhtml1/DTD/xhtml1-transitional.dtd
<html xmlns="http://www.w3.org/1999/xhtml">
<head>
<title>A for loop</title>
</head>
<body>
<script type="text/javascript" language="javascript">
document.write("<h3>a, b, c, = a*b + a*c + b*c<h3>")
B=28;
D = -160;
u = -300;
v = 300;
for (a = u; a \le v; a++)
        for (b = u; b \le v; b++)
                for (c = u; c \le v; c++)
if ((a+b+c==B) && (a*b*c==D))
```

The above html file gives the value of C in the cubic equation where B=28 and D=-160 $\,$

This is the output below:

```
a, b, c, = a*b + a*c + b*c

-5, 1, 32, = -133
-5, 32, 1, = -133
1, -5, 32, = -133
1, 32, -5, = -133
32, -5, 1, = -133
32, 1, -5, = -133
```

The above answer is divided by the value of $\mbox{\bf A}$ in the cubic equation to get the three solutions.

If A=2 then the solutions would be 1/2, -5/2 and 32/2